Upper Endoscopy EGD Prep Instructions

Please read the following instructions carefully

What is an **EGD**? (esophagogastroduodenoscopy) – An EGD is the visual examination of the upper intestinal tract esophagus, stomach, duodenum using a flexible scope that contains a light and miniature video camera on its tip; the scope is connected to a video screen for viewing.

WHAT TO DO BEFORE PROCEDURE: DYSPHAGIA (DIFFICULTY SWALLOWING)

• If you are currently taking blood thinners (for irregular heartbeat, atrial fibrillation, heart disease, stents, stroke, mini-stroke, mechanical heart valves, blood clots in the legs or lungs, or other reason) you may need to stop them prior to your procedure. Please contact the physician who prescribed the medication for instructions on how to stop them.

The following is not an all-inclusive list:

Plavix (clopidogrel bisulfate)	Effient (prasugrel)
Ticlid (ticlopidine hydrochloride)	Arixtra (fondaparinux)
Coumadin (warfarin sodium)	Brilinta (ticagrelor)
Pradaxa (dabigatran)	Lixiana/Savaysa (edoxaban)
Xarelto (rivaroxaban)	Lovenox (enoxaparin)
Eliquis (apixaban)	Fragmin (dalteparin)
Persantine (dipyridamole)	Zontivity (vorapaxar)
Pletal (cilostazol)	

- If you are taking full dose aspirin (325mg), please contact your prescribing physician to see if this can be decreased to low dose aspirin (81mg). **Do not stop low dose Aspirin (81mg)** Polyps can be removed in the presence of this lower dosage.
- If you have had a recent stress test or are scheduled to have one, please check with your physician to get a cardiac clearance. WE MUST HAVE RESULTS PRIOR TO PROCEDURE.
- If you are taking Adipex (Phentermine) or any GLP-1 Agonist (weight loss medication or for your diabetes) it is required to stop these medications 7 days prior to your procedure. Please contact your PCP regarding this if you are taking these medications for diabetes.
- Please refrain from marijuana use 12 hours prior to your procedure.

ONE DAY BEFORE PROCEDURE

- It is highly recommended to be on a **liquid diet** starting at **NOON** the day before your procedure
- Nothing to eat or drink after MIDNIGHT the night before your procedure

DAY OF PROCEDURE

- If you are diabetic you may check your blood sugar before coming in.
- **DO NOT EAT** or **DRINK** anything or your procedure may be cancelled.
- You <u>may take</u> all your medications with a sip of water except. blood sugar/diabetic medications.
 What to bring with you:
 - You cannot drive; you must be accompanied by an adult who must remain on the premises
 while you are having your procedure. Your procedure may be cancelled if you do not have a
 ride.
 - A list of your current medications including the dosage and the times you take them, along with any drug allergies.
 - Your driver's license and insurance cards