SIGMOIDOSCOPY PREPARATION INSTRUCTIONS

ONE WEEK PRIOR TO PROCEDURE:

- If you are currently taking blood thinners (for irregular heartbeat, atrial fibrillation, heart disease, stents, stroke, mini-stroke, mechanical heart valves, blood clots in the legs or lungs, or other reason) you may need to stop them prior to your procedure. Please notify us at 810-603-8400 prompt 1 and please contact the physician who prescribed the medication for instructions on how to stop them.
- If you are taking full dose aspirin (325mg), please contact your prescribing physician to see if
 this can be decreased to low dose aspirin (81mg). Do not stop low dose Aspirin (81mg).
 Polyps can be removed in the presence of this lower dosage.
- The following is not an all inclusive list.
- Plavix (clopidogrel bisulfate)
- Ticlid (ticlopidine hydrochloride)
- Coumadin (warfarin sodium)
- Pradaxa (dabigatran)
- Xarelto (rivaroxaban)
- Eliquis (apixaban)
- Persantine (dipyridamole)

- Effient (prasugrel)
- Arixtra (fondaparinux)
- Brilinta (ticagrelor)
- Lixiana/Savaysa (edoxaban)
- Lovenox (enoxaparin)
- Fragmin (dalteparin)
- Zontivity (vorapaxar)
- Pletal (cilostazol)
- If you are a diabetic, check with your family physician to see if you will need to adjust your insulin dosage.
- If you are taking Adipex (Phentermine) or any GLP-1 Agonist (weight loss medication or for your diabetes) it is required to stop these medications 7 days prior to your procedure. Please contact your PCP regarding this if you are taking these medications for diabetes.

ONE DAY BEFORE PROCEDURE:

- Light Dinner
- Nothing to eat or drink after MIDNIGHT the night before your procedure

ON THE DAY OF PROCEDURE:

- TWO (2) hours prior to your procedure you need to give yourself TWO (2) Fleets enemas before coming to the hospital
- If you are a diabetic you may check your blood sugar before coming in
- DO NOT EAT or DRINK anything or your procedure may be cancelled
- You <u>may take</u> all your medications with a sip of water EXCEPT: blood thinners, arthritis medications, weight loss medication, and blood sugar/diabetic medication
- What to bring with you:
- You are not allowed to drive, you MUST be accompanied by an adult who must remain on the
 premises while you're having your procedure. Your procedure may be cancelled if you do not
 have a ride. A list of your current medications including the dosage and time you take them,
 along with any drug allergies. Your driver's license and insurance cards.